

THE EFFECT OF ASYMMETRIC BEAMS IN THE WILKINSON MICROWAVE ANISOTROPY PROBE
EXPERIMENTI. K. WEHUS¹, L. ACKERMAN², H. K. ERIKSEN^{3,4}, AND N. E. GROENEBOOM^{3,4}*Draft version April 26, 2009*

ABSTRACT

We generate simulations of the CMB temperature field as observed by the WMAP satellite, taking into account the detailed shape of the asymmetric beams and scanning strategy of the experiment, and use these to re-estimate the WMAP beam transfer functions. This method avoids the need of artificially symmetrizing the beams, as done in the baseline WMAP approach, and instead measures the total convolution effect by direct simulation. We find noticeable differences with respect to the nominal transfer functions. For instance, the nominal V1 beam under-estimates the full beam convolution by $\sim 0.5\%$ at $\ell = 500$ and $\sim 1.0\%$ at $\ell = 800$. Similar differences are seen for other DA's. This in turn implies that the high- ℓ power spectrum is biased low by 1–2%, effectively tilting the spectrum slightly. Re-estimating cosmological parameters we find that the spectral index of scalar perturbations is $n_s = 0.969 \pm 0.014$ after correcting for this effect, corresponding to a positive shift of 0.3σ compared to the previously released WMAP results. Our CMB sky simulations are made publicly available, and can be used for general studies of asymmetric beam effects in the WMAP data.

Subject headings: cosmic microwave background — cosmology: observations — methods: statistical

1. INTRODUCTION

Without doubt, the angular CMB power spectrum is today our single most important source of cosmological information. Perhaps the most striking demonstration of this fact to date is the WMAP experiment, (Bennett et al. 2003; Hinshaw et al. 2007, 2009) which has allowed cosmologists to put unprecedented constraints on all main cosmological parameters, as well as ruling out vast regions of the possible model spaces. Similarly, in only a few years from now Planck will finally provide the definitive measurements of the temperature power spectrum, as well as polarization spectra with unprecedented accuracy. This will certainly lead to similar advances in our knowledge about the history of our universe.

Each of these experiments observes the CMB field by scanning the sky with an instrumental beam of finite resolution. This operation effectively corresponds to averaging over beam-sized angular scales, and is expressed technically either in pixel space by a convolution of the beam with the underlying sky, or in harmonic space by a multiplication of the two corresponding sets of harmonic expansion coefficients. For simplicity, the harmonic space expansion of the beam is typically expressed in terms of Legendre coefficients of an (azimuthally symmetric) effective beam response. This function is often called “the beam transfer function”, b_ℓ .

Before it is possible to make unbiased cosmological inferences based on the CMB power spectrum, it is of critical

importance to know the beam transfer function to high precision, as an error in the beam function translates into a direct bias in the estimated power spectrum. This in turn requires detailed knowledge about the beam response function on the sky for each experiment. For a full description of the WMAP beam estimation process and final model, see Page et al. (2003), Jarosik et al. (2007) and Hill et al. (2009).

The impact of asymmetric beams may also be important for applications other than power spectrum estimation. One example of special interest to us is the assessment of non-Gaussianity and violation of statistical isotropy. Specifically, Ackerman et al. (2007) considered a model based on violation of rotational invariance in the early universe, and derived explicit parametric expressions for the corresponding observational signature. Then, in a follow-up paper Groeneboom & Eriksen (2009) analysed the 5-year WMAP data with respect to this model and, most surprisingly, found a detection at the 3.8σ confidence level. Given that this was a most unexpected result, several questions concerning systematic errors in the WMAP data were considered, in particular those due to residual foregrounds, correlated noise and asymmetric beams. However, it was shown in the same paper that neither foregrounds nor correlated noise were viable explanations, while the question of asymmetric beams was left unanswered, due to lack of proper simulation machinery. This question provided our initial motivation for considering the problems studied in this paper.

The starting point for tackling the asymmetric beam problem for WMAP is a set of beam maps released by the WMAP team, two for each differencing assembly (DA), denoted A and B, respectively. These maps were derived by observing Jupiter for extended periods of time. Then, in order to derive the proper beam transfer functions, the WMAP team adopted a computationally fast and convenient approach: They first symmetrized the effective beam for each DA, collapsing the information in the A

Electronic address: i.k.wehus@fys.uio.no

Electronic address: lotty@caltech.edu

Electronic address: h.k.k.eriksen@astro.uio.no

Electronic address: leuat@irio.co.uk

¹ Department of Physics, University of Oslo, P.O. Box 1048 Blindern, N-0316 Oslo, Norway² California Institute of Technology, Pasadena, CA 91125³ Institute of Theoretical Astrophysics, University of Oslo, P.O. Box 1029 Blindern, N-0315 Oslo, Norway⁴ Centre of Mathematics for Applications, University of Oslo, P.O. Box 1053 Blindern, N-0316 Oslo, Norway

and B sides into one common function, and then computed the Legendre transform of the corresponding radial profile. However, for this to be an accurate approximation, one must on the one hand assume that the beams on the two sides are very similar, and on the other hand either assume that both beams are intrinsically circularly symmetric, or that all pixels on the sky are observed from all angles an equal number of times due to the scanning strategy. In reality none of these conditions are met, and one may therefore ask whether there might be any residual effect due to the combination of an asymmetric beam and anisotropic scanning in the WMAP beam functions.

This problem was addressed analytically by Hinshaw et al. (2007), who derived an approximate expression for the expected power spectrum bias due to asymmetric beams in the WMAP data. Their conclusion was that such effects were $\lesssim 1\%$ for the 3-year WMAP data.

In this paper, we revisit the question of asymmetric beams in WMAP with two main goals. First, we seek to estimate the effective beam transfer functions for each WMAP DA, taking into account the full details of the asymmetric beams and specifics of the WMAP scanning strategy by direct simulation. This way, we check whether the analytic approximations presented by Hinshaw et al. (2007) are valid. Second, we want to produce a set of high-fidelity simulated CMB sky maps, with beam properties as close as possible to those observed by WMAP, that can later be used for general studies of asymmetric beam effects in WMAP.

2. PIPELINE OVERVIEW

In this section we summarize the methods and algorithms used in this paper. Note that none of the individual steps described below are original to this paper, and only the main ideas will therefore be discussed in the following.

We begin by defining our notation. We will be estimating the product of the WMAP beam transfer function, b_ℓ , and pixel window, p_ℓ , by direct simulation. This product is denoted $\beta_\ell = b_\ell p_\ell$. Given this function, the combined effect on a sky map, $T(\hat{n})$, of convolution by an instrumental beam and averaging over finite-sized pixels may be approximated in harmonic space as

$$T(\hat{n}) = \sum_{\ell=0}^{\ell_{\max}} \sum_{m=-\ell}^{\ell} \beta_\ell a_{\ell m} Y_{\ell m}(\hat{n}), \quad (1)$$

where $Y_{\ell m}(\hat{n})$ are the usual spherical harmonics.

The angular power spectrum of T is given by

$$\hat{C}_\ell = \frac{1}{2\ell+1} \sum_{m=-\ell}^{\ell} \beta_\ell^2 |a_{\ell m}|^2, \quad (2)$$

while the power spectrum of the true underlying CMB map, $s(\hat{n})$, is

$$C_\ell = \frac{1}{2\ell+1} \sum_{m=-\ell}^{\ell} |a_{\ell m}|^2. \quad (3)$$

The effect of the beam convolution and pixel averaging on the power spectrum is therefore simply given by a multiplication with β_ℓ^2 .

The overall approach for estimating β_ℓ used in this paper may be summarized by the following steps: First, we simulate time-ordered data (TOD) for each DA, taking into account both the detailed beam maps of WMAP and the exact orientation of the spacecraft at each point in time. We then produce a sky map from this TOD. Finally we compute the square root of the ratio between the output and the input power spectra, which becomes our estimate of β_ℓ .

Note that in this paper we are only concerned with the effect of asymmetric beams, not other systematic effects such as instrumental noise. All following discussions will therefore assume noiseless observations.

2.1. Simulation of time-ordered data

Our first step is to simulate a reference CMB sky realization, s , given an angular temperature power spectrum, C_ℓ^{theory} . This can be achieved with a standard code such as “anafast”, which is available in the HEALPix⁵ software package. Note that this map should not be smoothed with either an instrumental beam or a pixel window; adding these effects is the task of the following pipeline. Explicitly, the input reference map should simply be pure spherical harmonic modes projected onto a set of pixel centers.

Next, we need to be able to convolve this map with a given beam map at arbitrary positions and orientations on the sphere. In this paper we do this by brute-force integration in pixel space. For an alternative fast Fourier space based approach to the same problem, see Wandelt & Górski (2001).

We define $\hat{p}$ to be a unit vector pointing towards the beam center, and specify its position on the sphere using longitude and co-latitude (ϕ, θ) . We further define ψ to be the angle between some fixed reference direction in the beam map and the local meridian. The value of the beam map at position $\hat{n} = (\phi', \theta')$, which in principle is non-zero over the full sky, is denoted $b(\phi', \theta'; \phi, \theta, \psi)$. With these definitions, the desired convolution may be written as

$$T(\phi, \theta, \psi) = \int_{4\pi} s(\phi', \theta') b(\phi', \theta'; \phi, \theta, \psi) d\Omega'. \quad (4)$$

Computationally speaking, we approximate this integral as a direct sum over HEALPix pixels, which all have equal area, with the product $s \cdot b$ being evaluated at HEALPix pixel centers. To make these calculations computationally feasible, we assume that the beam is zero beyond some distance from the beam center (ranging between 3.5 and 7° for the WMAP channels), and thus only include the main lobe in the following analysis. While the WMAP beam maps are provided as pixelized maps, we need to know the beam values at arbitrary positions (ie., HEALPix pixel centers). We solve this by computing a 2D spline for each beam map, enabling us to interpolate to arbitrary positions. For a review on one specific method for fast (and local) 2D spline evaluations, see Appendix B.

WMAP is a differential experiment, and measures at each point in time the difference between the signals received by two different detectors, denoted A and B. The

⁵ <http://healpix.jpl.nasa.gov>

full set of time-ordered WMAP data may therefore be written as

$$d_x(i) = T_x^A(i) - T_x^B(i), \quad (5)$$

where $x = \{K1, Ka1, Q1-2, V1-2, W1-4\}$ is a DA label, and i is a time index, and for each detector a shorthand for (ϕ, θ, ψ) . This equation may be written in the following matrix form,

$$\mathbf{d}_x = \mathbf{A}\mathbf{T}_x \quad (6)$$

where we have introduced an $N_{\text{tod}} \times N_{\text{pix}}$ pointing matrix $\mathbf{A}$. This matrix contains two numbers per row; 1 in the column hit by the center of beam A at time i , and -1 in the column hit by the center of beam B.

The remaining problem is to determine the position and orientation of each detector at each time step. This information has been made publicly available by the WMAP team on LAMBDA⁶, and consists of a large set of pointing files together with useful IDL routines for extracting the desired information.

2.2. Map making with differential data

For the map making step we adopt the algorithm developed by Wright et al. (1996), which was used in the 1- and 3-year WMAP pipelines (Hinshaw et al. 2003, 2007). Here we only summarize the essential algebra, and outline the algorithm.

Our goal is to establish an unbiased and, preferably, optimal estimate of the (smoothed) sky signal, $\hat{\mathbf{T}}$, given a set of differential TOD values, $\mathbf{d}$. For noiseless data, the maximum likelihood estimator is simply

$$\hat{\mathbf{T}} = (\mathbf{A}^t \mathbf{A})^{-1} \mathbf{A}^t \mathbf{d}. \quad (7)$$

For high-resolution sky maps, this equation involves an inverse of a large matrix and cannot be solved explicitly. Instead, one often resorts to iterative methods such as Conjugate Gradients, or, for differential data, the method developed by Wright et al. (1996).

We present the iterative differential map maker in a simple manner: Define $\mathbf{D}$ to be the diagonal matrix that counts the number of hits $N_{\text{obs}}(p)$ per pixel p on the diagonal, and a_i and b_i to be the pixels hit by side A and side B at time i , respectively. Suppose that we already have established some estimate for the solution, $\hat{\mathbf{T}}^j$. (Note that this can be zero.) Then the iterative scheme

$$\hat{\mathbf{T}}^{j+1} = \hat{\mathbf{T}}^j + \mathbf{D}^{-1} \mathbf{A}^t (\mathbf{d} - \mathbf{A} \hat{\mathbf{T}}^j) \quad (8)$$

will converge to the true solution: If $\hat{\mathbf{T}}^j = \mathbf{T}$, then $\mathbf{d} = \mathbf{A} \hat{\mathbf{T}}^j$, and the second term on the right hand side is zero. This algorithm is implemented by the following scheme:

$$\hat{T}_p^{j+1} = \frac{\sum_i \left(\delta_{p,a_i} \left[\hat{T}_{b_i}^j + d_i \right] + \delta_{p,b_i} \left[\hat{T}_{a_i}^j - d_i \right] \right)}{N_{\text{obs}}(p)}. \quad (9)$$

This algorithm was originally presented by Wright et al. (1996). The only new feature introduced here is the choice of starting point. In the original paper, Wright et al. (1996) initialized the iterations at the DMR dipole, since their test simulation included a CMB dipole term. However, for a given scanning strategy, there will often be some large-scale modes that




FIG. 1.— Top panel: Comparison between the transfer functions, β_ℓ , for a Gaussian beam of $20'$ FWHM. This was computed from a pixelized beam map and with the WMAP V1 scanning strategy (red line), and alternatively, by using the well-known analytic expression for the Legendre transform of a Gaussian beam (Equation 12) and isotropized HEALPix pixel window (black dashed line). The vertical dotted line indicates the multipole moment, ℓ_{hybrid} , at which $\beta_\ell = 0.15$. Bottom panel: The ratio between the transfer functions in the top panel. Note the excellent agreement up to $\ell \approx 800$, after which the differences in pixel window approximations becomes visible.




FIG. 2.— Comparison of the V1 WMAP beam transfer functions obtained from six (dashed line) and twelve (solid line) months of observations. Note that the behaviour at low and intermediate ℓ 's is similar, whereas the pixel window-induced increase is seen earlier in ℓ for the six month data set than for the twelve month data set.

are less well sampled than others. For instance, for the WMAP strategy $\ell = 5$ is more problematic than other modes (Hinshaw et al. 2003). This leads to slow convergence with the above scheme for this mode.

We therefore choose a different approach: Before solving for the high-resolution map by iterations, we solve Equation 7 by brute-force at low resolution. For the cases

⁶ <http://lambda.gsfc.nasa.gov>

TABLE 1
SUMMARY OF DA PARAMETERS

DA	FWHM ^a (arcmin)	Radius ^b (degrees)	N_{side}	ℓ_{max}	ℓ_{hybrid}	N_{samples} (10^8)	σ_N^c (μK)
K1	53	7.0	512	750	318	2.5	...
Ka1	40	5.5	512	850	411	2.5	...
Q1	31	5.0	512	1100	522	3.1	78.2
Q2	31	5.0	512	1100	515	3.1	74.2
V1	21	4.0	1024	1500	789	4.1	99.0
V2	21	4.0	1024	1500	779	4.1	88.2
W1	13	3.5	1024	1700	1164	6.2	143.8
W2	13	3.5	1024	1700	1148	6.2	159.7
W3	13	3.5	1024	1700	1162	6.2	168.5
W4	13	3.5	1024	1700	1169	6.2	164.4

^a Effective symmetrized beam size. ^b Radius used for pixelized beam convolutions. See Hill et al. (2009) for details. ^c Average full-sky RMS values evaluated at $N_{\text{side}} = 512$.

considered later in this paper, we choose a HEALPix resolution of $N_{\text{side}} = 16$ for this purpose. With 3072 pixels, about 30 seconds are needed to solve this system by singular value decomposition. (Note that the monopole is arbitrary for differential measurements, and one must therefore use an eigenvalue decomposition type algorithm to solve the system.) The improvement in convergence speed due to this choice of initial guess is explicitly demonstrated in Appendix A.

Our convergence criterion is chosen such that the RMS difference between two consecutive iterations must be less than $0.05 \mu\text{K}$. We have verified that this leads to errors of less than $0.1 \mu\text{K}$ in the final solution, of which far most is due to a residual dipole. This is typically achieved with 30–50 iterations, although some converge already after 20–30 iterations and a few after 70 or more iterations.

At first glance, the fact that the final residuals are as small as $0.1 \mu\text{K}$ for an RMS stopping criterion as large as $0.05 \mu\text{K}$ may seem surprising. However, this is explained by the fact that the iterative solution obtained by Equation 8 often alternates between high and low values about the true answer. This suggests that a further improvement to the algorithm may be possible: Faster convergence may perhaps be obtained by computing the average of two consecutive iterations, $\hat{\mathbf{T}} = (\hat{\mathbf{T}}^j + \hat{\mathbf{T}}^{j+1})/2$, as the map estimate for iteration $j + 2$. However, the computational resources spent during map making is by far sub-dominant compared to the TOD simulation, and we have therefore not yet implemented this step in our codes.

2.3. Estimation of hybrid beam transfer functions

As described in the introduction to this section, we estimate the transfer function by the square root of the ratio between the power spectra of the convolved map and the input map

$$\hat{\beta}_\ell^2 = \sqrt{\frac{\hat{C}_\ell}{C_\ell}}. \quad (10)$$

However, as noted above, this function describes both the effect from instrumental beam smoothing and averaging over pixels. In the present paper we are concerned mostly with the former of these, which has a stronger impact on large to intermediate scales. This is because the beam component is largely independent of total observation time, assuming at least one year of observations

for WMAP, whereas the pixel averaging component depends strongly on total observation time, or the average number of samples per pixel. The latter therefore evolves much more strongly with time than the former, as will be explicitly demonstrated later.

In order to provide transfer functions that are valid for long observation periods (e.g., 5 or 7 years of WMAP observations), we choose to construct a hybrid transfer function,

$$\hat{\beta}_\ell = \begin{cases} \sqrt{\frac{\hat{C}_\ell}{C_\ell}} & \text{for } \ell \leq \ell_{\text{hybrid}} \\ \sqrt{\frac{\hat{C}_{\ell_{\text{hybrid}}}}{C_{\ell_{\text{hybrid}}}}} \frac{b_{\ell_{\text{hybrid}}}^{\text{WMAP}}}{b_{\ell_{\text{hybrid}}}^{\text{WMAP}}} \frac{p_\ell}{p_{\ell_{\text{hybrid}}}} & \text{for } \ell > \ell_{\text{hybrid}} \end{cases}. \quad (11)$$

Here b_ℓ^{WMAP} is the nominal symmetrized transfer function published by the WMAP team, p_ℓ is the (uniformly averaged) HEALPix pixel window, and ℓ_{hybrid} is some transition multipole. In other words, we adopt our own direct estimate of the transfer function up to ℓ_{hybrid} , but the symmetrized, asymptotically uniform and properly scaled WMAP transfer function at higher multipoles.

Note that this issue is of minor importance in terms of cosmological interpretation, i.e., angular power spectrum and cosmological parameters, because the transition typically takes place in the noise dominated high- ℓ regime. The effect of the anisotropic pixel window is therefore largely suppressed. In the present paper, we therefore choose to focus on the beam dominated region, and leave a detailed study of the pixel window to a future paper. See Section 4 for a detailed discussion of this issue.

Finally, because we only generate a relatively small number of simulations in this paper, there is considerable Monte Carlo scatter in our estimated transfer functions on an ℓ -by- ℓ basis. To reduce this Monte Carlo noise, we smooth all transfer functions using the smooth spline formalism described by, e.g., Green & Silverman (1994).

3. DATA AND SIMULATIONS

All data products used in this study are provided by the WMAP team on LAMBDA as part of their 5-year data release. However, the calculations performed here are computationally extremely demanding, and we therefore include only roughly one year worth of data in our calculations. To be precise, we include the period between July 10th 2001 and August 2nd 2002, except for




FIG. 3.— Comparison between transfer functions derived in this paper (red lines) to the nominal WMAP transfer functions (black dashed lines). The transition multipole, ℓ_{hybrid} is marked by dotted vertical lines.

three days with missing data, for a total of 383 days⁷.

We consider all 10 WMAP DAs in our calculations, which are denoted, in order from low to high frequencies, K1 (23 GHz), Ka1 (33 GHz), Q1-2 (41 GHz), V1-2 (61 GHz) and W1-4 (94 GHz), respectively. Their resolutions range from 53' FWHM at K-band to 13' FWHM at W-band. Because of this large range in resolution, we specify the pixel resolution and harmonic space range for each case separately. For instance, K-band is pixelized at $N_{\text{side}} = 512$, and includes multipoles up to $\ell_{\text{max}} = 750$ (the highest multipole present in the transfer function provided by the WMAP team), while the W-band is pixelized at $N_{\text{side}} = 1024$, and includes multipoles up to $\ell_{\text{max}} = 1700$. A full summary of all relevant parameters for each DA is given in Table 1.

Note that the listed noise RMS values are only used for estimating the power spectrum weights in Section 6. For simplicity we have adopted the official RMS values for the foreground-reduced 5-year WMAP maps here, but note that there is a $\sim 1\%$ bias in some of these values (Groeneboom et al. 2009). However, this has no significant impact on the results presented in this paper.

The beam maps for each DA are provided in the form of pixelized maps, and separately for side A and B. Each beam map contains non-zero values inside a radius around the beam center which is specified for each DA. For instance, the K-band radius is 7° , and the W-band radius is 3.5° . When evaluating the convolution defined in Equation 4, we include all pixels inside this radius.

The pixel size of the beam maps is $2.4'$, which oversamples even the W-band beams. Based on these high-resolution maps we precompute all coefficients of the corresponding bi-cubic spline (see Appendix B), which allows us to very quickly interpolate at arbitrary positions in the beam map with high accuracy.

Each beam is normalized by convolving a map constant equal to 1 at 1000 random random positions and orientations, and demanding that the average of the resulting 1000 values equals unity. With the 2D spline interpolation scheme described in Appendix B the random uncertainties on the normalization due to beam position and orientation are $\sim 0.2\%$. For comparison, directly reading off pixel values from the beam maps without interpolation leads to variations in the normalization at the $\sim 2\%$ level.

For our base CMB reference sky set, we draw ten random Gaussian realizations from the the best-fit Λ CDM power spectrum derived from the 5-year WMAP data alone (Komatsu et al. 2009). These maps are generated at both ($N_{\text{side}} = 512$ and $N_{\text{side}} = 1024$ using the same seeds, and include neither an instrumental beam nor a pixel window; they are simply spherical harmonic modes projected onto the HEALPix pixel centers. All ten realizations are processed for all ten DAs, such that the resulting simulations may be used for multifrequency analysis, if so desired.

As noted above, the computational requirements for the analyses presented here are demanding. The CPU

time for processing a single W-band DA is ~ 4000 hours, and the total disk usage for the entire project is $\sim 1\text{TB}$. For comparison, the corresponding map making step requires ~ 60 CPU hours, and is thus completely subdominant the TOD simulation.

4. COMPARISON WITH ANALYTIC CASE

In order to test our pipeline and understand its outputs, we start by considering a perfect Gaussian beam. This case is treated in two different ways: First, we convolve a CMB realization directly in harmonic space (as defined by Equation 1) with a $\sigma_{\text{fwhm}} = 20'$ FWHM analytic Gaussian beam and the appropriate HEALPix pixel window, p_ℓ for $N_{\text{side}} = 1024$. The combined transfer function for this case reads

$$\beta_\ell^{\text{ref}} = e^{-\frac{1}{2}\ell(\ell+1)\sigma^2} p_\ell, \quad (12)$$

where $\sigma = \sigma_{\text{fwhm}}/\sqrt{8\ln 2}$, and σ_{fwhm} is expressed in radians.

Second, we map out a corresponding two-dimensional Gaussian in pixel space over a grid of $2.4'$ pixels, the same resolution as the WMAP beam maps. We then input this into our simulation pipeline together with the same CMB realization used for the analytic convolution, and with the V1 channel pointing sequence. From the resulting brute-force convolved map, we then obtain the effective transfer function, β_ℓ , as described in Section 2.3.

This function is plotted in the top panel of Figure 1, together with the product of the analytic Gaussian beam and the HEALPix pixel window. The ratio of the two effective functions is shown in the lower panel.

From this figure it is clear that the agreement between the two approaches is excellent up to $\ell \approx 800$. At higher ℓ 's, however, the ratio increases rapidly, indicating that the analytic approach smooths more than the brute-force approach. This is due to the different definitions of the pixel windows in the two cases: In the HEALPix case, the pixel window is defined as an effective average both over each pixel and over the full sky. In other words, it assumes that all points have been observed an equal (and infinite) number of times.

However, this is not the case for a real experiment which scans the sky for a finite length of time. As a consequence, each pixel is observed only a relatively small number of times, and this leads effectively to less smoothing. In the extreme case of only one observation per pixel, there would be no pixel averaging at all. Now, the average pixel window would formally equal unity. On the other hand, the random realization-specific uncertainties would be very large, and the pixel window as such would have zero predictive power.

In practice, one is well advised not to consider scales smaller than those that are properly oversampled by the scanning strategy. In this paper, we adopt the analytic case considered in this section to guide us in determining which scale that is. Explicitly, we conservatively demand that the effective beam transfer function must be greater than 0.15 in order to consider it to be properly oversampled, and therefore independent of scanning strategy. We adopt the corresponding multipole moment to be ℓ_{hybrid} , as defined in Section 2.3. Thus, the symmetrized WMAP beam and HEALPix pixel window are used at scales for which the beam amplitude drops below 0.15.

⁷ Our original intention was to include precisely one year of observations in our analysis, and therefore we processed 365 WMAP pointing files. However, we noticed after the calculations were completed that some of the pointing files contained slightly more than one day's worth of data, such that a total of 383 days was in fact included.




FIG. 4.— The ratio between the transfer functions derived in this paper and the nominal WMAP transfer functions for all DAs. Note that the DAs split into two main groups depending on focal plane position: The outer DAs, K1, Ka1 and Q1–2, all rise with ℓ , whereas the inner DAs, V1–2 and W1–4, decrease with ℓ . Note also the similarity between W1 and W4, between W2 and W3, and between V1 and V2.

Note that if we included more years of WMAP scanning in our calculations, we could increase ℓ_{hybrid} , since we would obtain more samples, and the WMAP pointings do not perfectly coincide from year to year. This is computationally very expensive, and we have therefore chosen to limit our analysis to the range described here.

As a direct illustration of this effect, we present in Figure 2 the transfer function ratios for the V1 WMAP DA with respect to the nominal WMAP transfer function, computed for both six months of observations and a full year of observations. The low and intermediate ℓ behaviour is the same for the two, but the pixel window effect becomes important earlier for the six month case than for the full year case.

5. THE EFFECT OF ASYMMETRIC BEAMS IN WMAP

We now present the main results obtained in this paper, namely the effective beam transfer functions for each WMAP DA, taking into account both the full asymmetric beam patterns and scanning strategy. These are shown in Figure 3 (red lines), and compared to the nominal WMAP transfer functions (dashed black lines). The vertical dotted line indicates ℓ_{hybrid} for each case.

Clearly, the differences between the two sets of results are relatively small, as no visual discrepancies are seen in this plot. However, in Figure 4 we plot the ratio between our transfer functions and the WMAP transfer functions for $\ell \leq \ell_{\text{hybrid}}$, and here we do see small but significant differences between the two sets of results.

First, we see that the ratios are essentially unity on the largest scales (smallest ℓ 's), before they start diverging either towards high or low values at some characteristic scale. There are two exceptions to this trend, namely W1 and W4, which start diverging essentially already at very low ℓ 's.

Next, the transfer functions split into two main groups: Our K1, Ka1 and Q1–2 transfer functions are slightly higher than the corresponding WMAP functions, while the V and W DAs are slightly lower.

Both of these general and qualitative remarks reflects the position of each DA in the WMAP focal plane (see Figure 6 of Jarosik et al. 2007 for an excellent visualization of the A side beams): K1, Ka1 and Q1–2 are positioned the furthest away from the optical axis, while V1–2 and W1–4 are the closest. Similarly, W1 and W4 are positioned lower in elevation, and generally have more sub-structure, than W2 and W3.

However, it should be emphasized that the overall differences are generally small, typically less than 2% at $\ell \leq \ell_{\text{hybrid}}$. Further, these differences are only significant (again, with the exception of W1 and W4) in the intermediate- and high- ℓ ranges.

To build up some intuitive understanding of the spatial variations caused by the asymmetric WMAP beams, we show in Figure 5 the difference between the fully asymmetrically convolved map and the corresponding map convolved with the symmetrized transfer function directly in harmonic space for one of the V1 simulations. Thus, the two convolved maps have identical power spectra, but slightly different phases. The top panel shows the full-sky difference map with a temperature scale of $\pm 5\mu\text{K}$. The lower panels show two selected $15^\circ \times 15^\circ$ regions centered on the north ecliptic pole (NEP; top row) and the Galactic center (GC; bottom row), respectively.

The left column shows the actual temperature map convolved with the asymmetric beam, and the right column shows the same differences as in the full-sky plot.

The first striking feature seen in this map is that the differences are clearly larger in the ecliptic plane than around the ecliptic poles. This is due to the WMAP scanning strategy, which leads to a larger number of observations per pixels around the poles, and also with a greater range of beam orientations. Next, it is difficult to spot any single unambiguous and well-defined correlation between the convolved and the difference maps. Clearly, there are similar morphological structures in the two, but the sign of the correlations appears to vary. Third, we see a clear tendency of diagonal striping in the GC plot, which corresponds to correlations along ecliptic meridians and lines of constant latitude. (Note that these plots are shown in Galactic coordinates, while the WMAP scanning strategy is nearly azimuthally symmetric in ecliptic coordinates.)

In the next section, we consider the impact of the asymmetric beams on cosmological parameters. However, before concluding this section we make a comment concerning an outstanding issue regarding the 3-year WMAP power spectra first noted by Eriksen et al. (2007). They pointed out the presence of a 3σ amplitude discrepancy between the V- and W-band power spectra (Figure 5 of Eriksen et al. 2007). Specifically, the V-band spectrum was biased low compared to the W-band spectrum between $\ell = 300$ and 600 by $\sim 80\mu\text{K}^2$. Hufenberger et al. (2006) later showed that $\sim 30\mu\text{K}^2$ of this discrepancy could be attributed to over-estimation of point source power in the 3-year WMAP spectrum analysis, and this was subsequently confirmed and corrected by Hinshaw et al. (2007). Still, about $50\mu\text{K}^2$ of this difference remained, which was statistically significant at $\sim 2\sigma$.

Eriksen et al. (2007) proposed that this difference might be due to errors in the beam transfer functions caused by asymmetric beams. Given the new results presented in this paper, we are now in a position to consider this issue more quantitatively. The relevant question is then whether the WMAP V-band transfer functions are systematically biased high compared to the W-band functions. At first glance, one may get this impression from the plots shown in Figure 4: The V-band ratios both drop noticeably from $\ell = 300$ (decreasing nearly linearly from -0.2 to -0.7%), whereas W2 and W3 are slightly high in the same range, at about +0.1 to +0.2%. On the other hand, W1 and W4 are even lower than the V-band functions, at -0.4 to -0.6%. The net difference is therefore not more than a few tenths of a percent, which corresponds to $\sim 10\mu\text{K}^2$ in the power spectrum. Thus, it is possible that this effect may contribute somewhat to the power spectrum discrepancy between V- and W-band, but it does not seem to fully explain the difference.

6. IMPACT ON COSMOLOGICAL PARAMETERS

We now assess the impact of asymmetric beams in WMAP on cosmological parameters. We do this by modifying the co-added 5-year WMAP temperature power spectrum (Nolta et al. 2009) provided with the WMAP likelihood code Dunkley et al. (2009); Komatsu et al. (2009) according to the transfer function ratios shown in Figure 4, and run CosmoMC (Lewis & Bridle 2002)




FIG. 5.— Top panel: Difference between a V1 simulation convolved with the full asymmetric beam and the same realization convolved with the corresponding symmetrized transfer function. The two maps have identical power spectrum but different phases. Note that larger differences are observed along the ecliptic plane, where the density of observations is lower than towards the ecliptic poles, and the cross-linking is also weaker. Bottom panels: Zoom-in on two regions, the north ecliptic pole (NEP; top row) and the Galactic center (GC; bottom row). Left column shows the map convolved with an asymmetric beam, and right column shows the same difference as in the top panel.




FIG. 6.— Total correction to the 5-year co-added WMAP temperature power spectrum due to asymmetric beams. Note the transition between high and low signal-to-noise weighting schemes at $\ell = 500$, and also the manually capped amplitude at $\ell > 750$. The latter is imposed in order to be conservative in the very high- ℓ regime, where the transfer functions are sensitive to pixel window effects.

to estimate the resulting parameters. Only a simple 6-parameter Λ CDM model is considered in this paper. For comparison, we also run the code with the nominal WMAP spectrum as input, so that we can directly estimate the impact of asymmetric beams with everything else held fixed.

Unfortunately, the individual cross-spectra for each pair of DAs have not yet been published by the WMAP team, but only the total co-added spectrum. We must therefore make a few approximations in order to apply the proper beam corrections to the full spectrum. First, let σ_n^i denote the white noise level of DA i (see Table 1), β_ℓ^i the transfer function estimate derived in this paper, and let $\beta_\ell^{i,\text{WMAP}}$ be the nominal WMAP transfer function. Finally, define

$$\delta_\ell^i = \frac{\beta_\ell^i}{\beta_\ell^{i,\text{WMAP}}} - 1 \quad (13)$$

to be the fractional difference between the two.

Next, the WMAP team uses the MASTER pseudo-spectrum algorithm (Hivon et al. 2002) for power spectrum estimation (Hinshaw et al. 2003, 2007; Nolte et al. 2009), which quickly produces good estimates at high ℓ 's. However, this method is not a maximum-likelihood estimator, and it does not yield optimal error bars. To improve on this, the WMAP applies different pixel weights in different multipole regions: At low ℓ 's, where the sky maps are signal dominated, they apply equal weights to all pixels, while at high ℓ 's, where the maps are noise dominated, they apply inverse noise variance pixels weights. These weights are then taken into account when co-adding the cross-spectra obtained from all possible DA pairs (but excluding auto-correlations). The transition is made at $\ell = 500$.

The beam-convolved (but noiseless) power spectrum $\tilde{C}_\ell^{ij}$ observed by a given DA pair, i and j , may be written as $\tilde{C}_\ell^{ij} = \beta_\ell^i \beta_\ell^j C_\ell$, where C_ℓ is the true power spectrum of our sky, and β_ℓ^i is the true transfer function for DA i . The noise amplitude of the same spectrum is proportional to $\sigma_n^i \sigma_n^j / \beta_\ell^i \beta_\ell^j$. The inverse noise variance weight of this

cross-spectrum is therefore approximately

$$w_{ij} = \frac{\frac{\beta_\ell^i \beta_\ell^j}{\sigma_n^i \sigma_n^j}}{\sum_{i' < j'} \frac{\beta_\ell^{i'} \beta_\ell^{j'}}{\sigma_n^{i'} \sigma_n^{j'}}}, \quad (14)$$

where the sum runs over all N different pairs of cross-spectra. (Note that this is only an approximation to the exact expression, because other effects also enter the full calculations. One important example is the sky cut, which couples different ℓ modes, and is taken into account through a coupling matrix. Such effects are not included in the analysis presented here.

Pulling all of this together, the appropriately co-added power spectrum provided by WMAP should ideally read

$$\hat{C}_\ell = \begin{cases} \frac{1}{N} \sum_{i < j} \frac{\tilde{C}_\ell^{ij}}{\beta_\ell^i \beta_\ell^j} & \text{for } \ell \leq 500 \\ \sum_{i < j} w_{ij} \frac{\tilde{C}_\ell^{ij}}{\beta_\ell^i \beta_\ell^j} & \text{for } \ell > 500 \end{cases} \quad (15)$$

However, the spectrum that in fact is provided by WMAP is Equation 15 evaluated for $\beta_\ell = \beta_\ell^{\text{WMAP}}$, which, according to our calculations, is slightly biased. To obtain the appropriate correction factor, $\alpha_\ell = \hat{C}_\ell / \hat{C}_\ell^{\text{WMAP}}$, for each ℓ , we therefore set $\beta_\ell = \beta_\ell^{\text{WMAP}} (1 + \delta_\ell)$ in Equation 15, and expand to first order in δ_ℓ . Doing this, we find that

$$\alpha_\ell = \begin{cases} 1 - \frac{1}{N} \sum_{i < j} (\delta_\ell^i + \delta_\ell^j) & \text{for } \ell \leq 500 \\ 1 - \sum_{i < j} w_{ij}^{\text{WMAP}} (\delta_\ell^i + \delta_\ell^j) & \text{for } \ell > 500 \end{cases} \quad (16)$$

where w_{ij}^{WMAP} is the expression given in Equation 14 evaluated with β_ℓ^{WMAP} .

This function is plotted in Figure 6. Note, however, that we have capped the function by hand at $\ell = 750$ to be conservative, considering that our V-band transfer functions do not have support all the way to the maximum multipole used in the WMAP likelihood code, $\ell_{\text{max}} = 1000$.

The results from the corresponding CosmoMC analyses are tabulated in Table 2 in terms of marginal means and standard deviations, and plots of the marginal distributions are shown in Figure 7. Here we see that there are small but noticeable shift in several parameters. For example, there is a positive shift of 0.4σ in the amplitude of scalar perturbations, A_s , and 0.3σ in the spectral index of scalar perturbations. Although relatively modest, these shifts are certainly large enough that they should be taken properly into account.

7. CONCLUSIONS

This paper has two main goals. First, we wanted to generate a set of WMAP-like simulations that fully take into account the asymmetric beams and anisotropic scanning pattern of the WMAP satellite. Such simulations are extremely valuable for understanding the impact of beam asymmetries on various statistical estimators and models. One example of such, which indeed provided us with the initial motivation for studying this issue, is the anisotropic universe model presented by Ackerman et al. (2007), and later considered in detail with respect to the WMAP data by Groeneboom & Eriksen (2009). The result from that analysis was a tentative detection of violation of rotational invariance in the early universe, or




FIG. 7.— Marginal distributions for cosmological parameters derived with (dashed lines) and without (solid lines) correction for asymmetric beams.

TABLE 2
COSMOLOGICAL PARAMETERS

Parameter	Nominal WMAP	Corrected beams	Shift in σ
$\Omega_b h^2$	0.0228 ± 0.0006	0.0228 ± 0.0006	0.1
$\Omega_{\text{cdm}} h^2$	0.109 ± 0.0006	0.112 ± 0.006	0.4
$\log(10^{10} A_s)$	3.064 ± 0.042	3.079 ± 0.042	0.4
τ	0.089 ± 0.017	0.089 ± 0.017	0.00
h	0.722 ± 0.027	0.716 ± 0.026	-0.3
n_s	0.965 ± 0.014	0.969 ± 0.014	0.3

NOTE. — Comparison of cosmological parameters derived from the nominal WMAP power spectrum (second column) and from the power spectrum corrected for asymmetric beams (third column). The rightmost column shows the relative shift between the two in units of σ .

some other effect with similar observational signatures, at the 3.8σ confidence level. It was shown that neither foregrounds nor correlated noise could generate this signal, but the question of asymmetric beams was left unanswered. This issue will now be revisited in an upcoming paper, using the simulations generated here.

The second goal of the paper was to assess the impact of beam asymmetries on the WMAP power spectrum and cosmological parameters. We did this by comparing the power spectrum of the full beam convolved simulations with the power spectrum of the input realizations, thereby providing a direct estimate of the effective beam transfer functions. Doing so, we found differences at the 1–2% level in all differencing assemblies at intermediate and high ℓ 's with respect to the nominal WMAP transfer functions.

A similar analysis was performed for the 3-year WMAP data release by Hinshaw et al. (2007), who approach the problem from an analytical point of view. However, at that time only the A-side beams were available (Hill et al. 2009), and they therefore assumed identical beams on both the A and B sides. With this data,

they concluded that the impact of beam asymmetries was $\lesssim 1\%$ everywhere below $\ell = 1000$ for the V- and W-band DAs. For comparison, we find that there is a $\sim 1\%$ bias already at $\ell = 600$ for the combined co-added temperature power spectrum, and increasing rapidly to $\sim 1.5\%$ at $\ell = 750$. It is not unlikely that this trend may continue further in ℓ , but to answer that question considerably more computational resources are required. Nevertheless, the two analyses appear to be in reasonable agreement with each other, especially considering the fact that we take into account the full beam maps of both the A and B sides.

As far as cosmological parameters go, the impact of asymmetric beams appears to be small but noticeable. Specifically, we find shifts of 0.4σ in the amplitude of scalar perturbations, A_s , and the physical density of cold dark matter, $\Omega_{\text{cdm}} h^2$, and 0.3σ in the spectral index of scalar perturbations, n_s . While these shifts are relatively modest, they are of the same order of magnitude or larger than, say, marginalization over the Sunyaev-Zeldovich effect (Spergel et al. 2007) or unresolved point sources (Nolta et al. 2009), which indeed are taken into account.

One outstanding question that still remains is the impact of the anisotropic effective pixel window. As explicitly demonstrated in this paper, the difference between the isotropized HEALPix pixel window and the actual WMAP V1 scanning induced pixel window becomes visible at $\ell \sim 900$ for one year of WMAP observations. Of course, this is well within the noise-dominated regime for the WMAP data, and unlikely to have any major impact on cosmological results, but we believe that a proper understanding of this issue, both with respect to WMAP and Planck, is warranted, and intend to revisit this issue in a separate study.

The simulations described in this paper may be downloaded from IKW's homepage⁸.

The authors thank Ned Wright for very useful and

⁸ http://www.fys.uio.no/~ingunnkw/WMAP5_beams

stimulating discussions, which eventually led to the conception of this project. We also thank Kris Górski, Sanjit Gupta, Gary Hinshaw and Graca Rocha for interesting discussions. LA thanks the Institute of Theoretical Astrophysics, Oslo, for its generous hospitality during her visit to Norway. IKW, HKE and NEG acknowledge financial support from the Research Council of Norway. The computations presented in this paper were carried out on Titan, a cluster owned and maintained by the University of Oslo and NOTUR. Some of the results in this paper have been derived using the HEALPix (Górski et al. 2005) software and analysis package. We acknowledge use of the Legacy Archive for Microwave Background Data Analysis (LAMBDA). Support for LAMBDA is provided by the NASA Office of Space Science.

APPENDIX

CONVERGENCE OF THE DIFFERENTIAL MAP MAKER

As described in Section 2.2, we introduce one new step to the differential map making algorithm presented by Wright et al. (1996): We initialize the iterations at the exact solution of Equation 7 evaluated at low resolution, which in this paper is taken to be $N_{\text{side}} = 16$, with 3072 pixels.

To demonstrate the improvement in convergence due to this choice of initialization, we revisit the analytic case considered in Section 4, which compared the results from our simulation pipeline with an exact analytic case, but taking into account the actual WMAP scanning strategy.

In Figure 8 we show a set of difference maps taken between the intermediate solutions produced by the differential map maker and the analytic and isotropic map solution. From top to bottom, the panels show the residuals after 2, 5 and 10 iterations, and at the bottom, the final converged solutions. The left panel shows the series obtained when initializing the search at the low-resolution solution, while the right panel shows the series when initializing at zero. Convergence was achieved respectively after 67 and 123 iterations in the two cases.

Note that the WMAP team initializes their search at the CMB dipole, which is the dominant component in their data set. However, this is in our setting equivalent to initializing at zero, since our simulation does not include a dipole.

Taking the difference between the two final solutions, we have verified that the peak-to-peak residuals in the two maps are less than $0.1 \mu\text{K}$, of which essentially all is concentrated in a single dipole component. The solution is thus independent of initialization, and the only difference lies in computational speed.

Finally, note that even though the two maps are internally indistinguishable, they are both quite different from the isotropic reference map. To be precise, the RMS difference between the derived maps and the isotropic reference map is $0.91 \mu\text{K}$, with a spatial pattern similar to the overall WMAP scanning pattern.

The cause of these residuals is once again the differences in the treatment of the effective pixel windows: The HEALPix pixel window is computed by uniformly averaging over the full sky, whereas the simulation pipeline takes into account the actual pointing directions of the satellite. Sub-pixel variations in the CMB sky therefore leads to significant differences in the two estimates on small scales. The effect of such pixel window variations on the 5-year WMAP power spectrum will be considered in a future paper.

FAST 2D SPLINE EVALUATION

The heart of the simulation pipeline described in Section 2 is the real-space convolution algorithm defined by Equation 4. For this operation to be computationally feasible we have to be able to evaluate the beam response function quickly at any position. The real beam maps, however, are provided to us in the form of two-dimensional pixelized images with relatively coarse resolution. It is therefore necessary to establish a fast and accurate interpolation scheme.

We adopt a bicubic spline for this purpose, and review here one specific implementation of this concept. Note that most of the following is standard textbook material (e.g., Press 2002), and is included here only for easy reference.

Suppose we are given some tabulated two-dimensional function $f(x, y)$ over a regular grid, and want to interpolate at arbitrary positions (x_0, y_0) within this grid. One particularly appealing approach for doing so are by means of bicubic splines, which are bi-cubic polynomials in x and y ,

$$p(x, y) = \sum_{i=0}^3 \sum_{j=0}^3 a_{ij} x^i y^j. \quad (\text{B1})$$

The coefficients a_{ij} are defined separately for each grid cell, and our task is to compute these given the tabulated function $f(x, y)$. Note that once we have these coefficients, any spline evaluation will be very fast, since it essentially




FIG. 8.— Comparison of convergence of the differential map maker for two different choices of initialization. The left column shows the snapshots from the series obtained with initializing at a solution obtained by brute-force evaluation at low resolution, while the right column shows the series obtained when initializing at zero. Each plot is a difference map between the current solution for a data set including asymmetric beams and real scanning strategy and the corresponding map convolved with the analytic Gaussian beam and isotropic HEALPix pixel window. The bottom row shows the final solutions obtained in the two cases, which were obtained after 67 and 123 iterations, respectively. These final maps are identical up to a $\sim 0.1\mu\text{K}$ dipole.

amounts to performing a vector-matrix-vector multiplication with a 4×4 matrix.

Let us first consider a cell defined over the unit square, having corners $(x, y) = (0, 0)$, $(1, 0)$, $(0, 1)$ and $(1, 1)$. (Note that this assumption does not imply any restriction of the problem, since any grid cell in a regular grid may be linearly transformed into the unit square.) Assume also that we know the function values $f(x, y)$ and the first- and second-order derivatives, $f_x(x, y)$, $f_y(x, y)$ and $f_{xy}(x, y)$ at all four corners. (Here subscript x denotes derivatives, $f_x = df/dx$.) The

coefficients of the bicubic spline are then defined such that both the function values and the derivatives match,

$$f(x, y) = p(x, y) \quad (\text{B2})$$

$$f_x(x, y) = p_x(x, y) \quad (\text{B3})$$

$$f_y(x, y) = p_y(x, y) \quad (\text{B4})$$

$$f_{xy}(x, y) = p_{xy}(x, y), \quad (\text{B5})$$

at all four corners. With four equations for each of four corners, there is a total of 16 independent equations for the 16 independent spline coefficients, a_{ij} , and the spline is therefore uniquely defined.

Writing out Equation B5 explicitly, we obtain the following set of linear equations,

$$\left. \begin{aligned} f(0, 0) &= p(0, 0) = a_{00} \\ f(1, 0) &= p(1, 0) = a_{00} + a_{10} + a_{20} + a_{30} \\ f(0, 1) &= p(0, 1) = a_{00} + a_{01} + a_{02} + a_{03} \\ f(1, 1) &= p(1, 1) = \sum_{i=0}^3 \sum_{j=0}^3 a_{ij} \end{aligned} \right\} \text{ Matching function values} \quad (\text{B6})$$

$$\left. \begin{aligned} f_x(0, 0) &= p_x(0, 0) = a_{10} \\ f_x(1, 0) &= p_x(1, 0) = a_{10} + 2a_{20} + 3a_{30} \\ f_x(0, 1) &= p_x(0, 1) = a_{10} + a_{11} + a_{12} + a_{13} \\ f_x(1, 1) &= p_x(1, 1) = \sum_{i=0}^3 \sum_{j=0}^3 i a_{ij} \end{aligned} \right\} \text{ Matching first-order } x\text{-direction derivatives} \quad (\text{B7})$$

$$\left. \begin{aligned} f_y(0, 0) &= p_y(0, 0) = a_{01} \\ f_y(1, 0) &= p_y(1, 0) = a_{01} + a_{11} + a_{21} + a_{31} \\ f_y(0, 1) &= p_y(0, 1) = a_{01} + 2a_{02} + 3a_{03} \\ f_y(1, 1) &= p_y(1, 1) = \sum_{i=0}^3 \sum_{j=0}^3 j a_{ij} \end{aligned} \right\} \text{ Matching first-order } y\text{-direction derivatives} \quad (\text{B8})$$

$$\left. \begin{aligned} f_{xy}(0, 0) &= p_{xy}(0, 0) = a_{11} \\ f_{xy}(1, 0) &= p_{xy}(1, 0) = a_{11} + 2a_{21} + 3a_{31} \\ f_{xy}(0, 1) &= p_{xy}(0, 1) = a_{11} + 2a_{12} + 3a_{13} \\ f_{xy}(1, 1) &= p_{xy}(1, 1) = \sum_{i=0}^3 \sum_{j=0}^3 ij a_{ij} \end{aligned} \right\} \text{ Matching second-order } xy \text{ cross-derivatives} \quad (\text{B9})$$

The remaining problem is then to estimate the first- and second-order derivatives at all grid points, and several approaches may be used for this. We adopt a spline based method for this step as well.

First, we compute a standard one-dimensional natural spline along all x and y coordinate lines, using standard methods. The result from this operation is the set of all pure second-order derivatives, $f_{xx}(x, y)$ and $f_{yy}(x, y)$, at each grid point.

Because a one-dimensional spline is also a simple cubic polynomial, and therefore only has four free coefficients, it is sufficient to know the function values and second-order derivatives at all grid corners to uniquely specify every coefficient. As a consequence of this, the first-order derivative along the x -direction at grid point (x_i, y_j) is uniquely given by (Press 2002)

$$f_x(x_i, y_j) = \frac{f(x_{i+1}, y_j) - f(x_i, y_j)}{h_x} - \frac{1}{3} h_x f_{xx}(x_i, y_j). \quad (\text{B10})$$

Here $h_x = x_{i+1} - x_i$ is the x -direction grid cell size for the current cell. An equivalent expression obviously holds for the y -direction derivatives.

Finally, to estimate the second-order cross-derivatives, $f_{xy}(x, y)$, the above process is repeated such that y -derivatives are computed from $f_x(x, y)$ splines for all y -direction coordinate lines. Thus, all required derivatives may be obtained by performing $m + 2n$ one-dimensional spline computations, where m is the number of grid cells in x -direction, and n is the number of grid cells in y -direction.

REFERENCES

- Ackerman, L., Carroll, S. M., & Wise, M. B. 2007, *Phys. Rev. D*, 75, 083502
- Bennett, C. L., et al. 2003, *ApJS*, 148, 1
- Dunkley, J., et al. 2009, *ApJS*, 180, 306
- Eriksen, H. K., et al. 2007, *ApJ*, 656, 641
- Górski, K. M., Hivon, E., Banday, A. J., Wandelt, B. D., Hansen, F. K., Reinecke, M., Bartelman, M. 2005, *ApJ*, 622, 759
- Green, P. J., & Silverman, B. W. 1994, *Non-Parametric Regression and Generalized Linear Models*, Chapman and Hall, 1994
- Groeneboom, N. E., & Eriksen, H. K. 2009, *ApJ*, 690, 1807
- Groeneboom, N. E., Eriksen, H. K., Gorski, K., Huey, G., Jewell, J., & Wandelt, B. 2009, *arXiv:0904.2554*
- Hill, R. S., et al. 2009, *ApJS*, 180, 246
- Hinshaw, G., et al. 2003, *ApJS*, 148, 63
- Hinshaw, G., et al. 2007, *ApJS*, 170, 288
- Hinshaw, G., et al. 2009, *ApJS*, 180, 225
- Hivon, E., Górski, K. M., Netterfield, C. B., Crill, B. P., Prunet, S., & Hansen, F. 2002, *ApJ*, 567, 2
- Huffenberger, K. M., Eriksen, H. K., & Hansen, F. K. 2006, *ApJ*, 651, L81
- Jarosik, N., et al. 2007, *ApJS*, 170, 263
- Komatsu, E., et al. 2009, *ApJS*, 180, 330
- Lewis, A., & Bridle, S. 2002, *Phys. Rev. D*, 66, 103511
- Nolta, M. R., et al. 2009, *ApJS*, 180, 296
- Page, L., et al. 2003, *ApJS*, 148, 39
- Press, W. H. 2002, "Numerical recipes in C++ : the art of scientific computing", Cambridge University Press, ISBN: 0521750334
- Spergel, D. N., et al. 2007, *ApJS*, 170, 377
- Wandelt, B. D., & Górski, K. M. 2001, *Phys. Rev. D*, 63, 123002
- Wright, E. L., Hinshaw, G., & Bennett, C. L. 1996, *ApJ*, 458, L53